



Sharif University of Technology

Scientia Iranica

Transactions C: Chemistry and Chemical Engineering

www.sciencedirect.com

One-pot synthesis of 1, 2, 4, 5-tetrasubstituted imidazoles promoted by nano-TiCl₄.SiO₂

B.F. Mirjalili^{a,*}, A.H. Bamoniri^b, L. Zamani^a^a Department of Chemistry, College of Science, Yazd University, Yazd, P.O. Box 8915813149, Iran^b Department of Organic Chemistry, Faculty of Chemistry, University of Kashan, Kashan, P.O. Box 51167, Iran

Received 13 August 2011; revised 25 October 2011; accepted 17 December 2011

KEYWORDS

1, 2, 4, 5-tetrasubstituted imidazoles;
Nano-TiCl₄.SiO₂;
Benzil;
One-pot synthesis;
Solvent free condition;
Supported Lewis acid.

Abstract Nano-TiCl₄.SiO₂ is an efficient, readily available and reusable catalyst for the synthesis of 1, 2, 4, 5-tetrasubstituted imidazoles using benzil, aromatic aldehyde and an amine in the presence of ammonium acetate. The one-pot procedure is very simple with good to excellent yields.

© 2012 Sharif University of Technology. Production and hosting by Elsevier B.V.

Open access under [CC BY license](http://creativecommons.org/licenses/by/4.0/).

1. Introduction

The imidazole core is present in many molecules with biological activity [1], such as anti-allergy [2], anti-inflammatory [3], analgesic, antifungal, antimycotic, antibiotic, antiulcerative, antibacterial and antitumor [4]. In addition, many of the substituted diaryl imidazoles are known as potential inhibitors of the p38 MAP kinase [5]. Appropriately, substituted imidazoles are extensively used as glucagon receptors [6] and CB1 cannabinoid receptor antagonists [7], modulators of P-glycoprotein (P-gp)-mediated multidrug resistance (MDR) [8], antibacterial [9] and antitumor [10] agents.

Recently, the synthesis of 1, 2, 4, 5-tetrasubstituted imidazoles has been catalyzed by silica gel or Zeolite HY [11], silica gel/NaHSO₄ [12], I₂ [13], K₅CoW₁₂O₄₀.3H₂O [14], heteropolyacids [15], HClO₄-SiO₂ [16], InCl₃.3H₂O [17], ZrCl₄ [18],

BF₃.SiO₂ [19], DABCO [20], PEG-400 [21] and silica-bonded propylpiperazine *N*-sulfamic acid (SBPPSA) [22].

TiCl₄.SiO₂, as a powerful Lewis acid, is a liquid which is highly volatile, corrosive and difficult to handle. It hydrolyses to produce HCl in the presence of moisture. Silica-supported TiCl₄ is a mild solid Lewis acid that promotes acidic catalyzed organic reactions. This catalyst does not need special precautions for preparation, handling or storage, and it can be stored at an ambient temperature for months without losing its catalytic activity. This catalyst was previously applied for the production of polymers [23].

In continuation of our investigation into the application of solid acids in organic synthesis [19,24–26], we have applied Nano-TiCl₄.SiO₂ as an efficient catalyst for synthesis of 1, 2, 4, 5-triphenylimidazoles via the reaction of benzil, aldehydes, amine and ammonium acetate.

2. Results and discussion

Nano silica supported titanium chloride (nano-TiCl₄.SiO₂), as an efficient acidic catalyst, was prepared via a reaction of nano SiO₂ with TiCl₄. The dimensions of nanoparticles were observed with SEM. The particle sizes of the commercial silica gel and synthesized nano-TiCl₄.SiO₂ were about 24 and 41 nm, respectively (Figure 1).




* Corresponding author.

E-mail address: fmirjalili@gmail.com (B.F. Mirjalili).

Peer review under responsibility of Sharif University of Technology.



Production and hosting by Elsevier

Figure 1: SEM photograph of (a) nano-SiO₂, and (b) nano-TiCl₄.SiO₂.Figure 2: FT-IR spectrum of (a) nano-SiO₂, (b) nano-TiCl₄.SiO₂, and (c) TiCl₄.Figure 3: Suggested structure for nano-TiCl₄.SiO₂.

For identification of the structure of nano TiCl₄.SiO₂, we have studied the IR spectra of nano-SiO₂, nano TiCl₄.SiO₂ and TiCl₄ (Figure 2). In all of the spectra, very broad OH stretching bands are observed, with strong intermolecular hydrogen bonding. In FT-IR spectra of nano TiCl₄.SiO₂ and SiO₂, the absorption bands for Si-OH and Si-O-Si appear at ~700 and ~1100 cm⁻¹, respectively. The absorption band of Ti-Cl appears at 1600 cm⁻¹ in the TiCl₄ spectrum. In the FT-IR spectrum of nano-TiCl₄.SiO₂, the O-Ti-Cl, Si-OH and Si-O-Si absorption bands are observed at 900, 700 and 1100 cm⁻¹, respectively [27,28]. We have suggested the following structure for nano-TiCl₄.SiO₂ (Figure 3).

Because the nano-TiCl₄.SiO₂ produces HCl in water, we have determined the concentration of the obtained acidic aqueous solution by titration. We have found that 0.1 g of catalyst produced a 0.0064 M acidic aqueous solution. For determination of the loading amount of Ti on 0.1 g of nano-TiCl₄.SiO₂, we have extracted Ti from the catalyst using an EDTA solution (0.004 M) at pH of 4. The standard solutions of titanium (IV) were prepared by dissolving a proper amount of TiCl₄ in acetone. Determination of the extracted Ti was carried out by an atomic absorption spectrometer with a hollow cathode lamp, at a wavelength of 346 nm, using C₂H₂/N₂O flame. The calibration equation for titanium was found as $A = 0.012C - 0.1895$





Figure 4: Synthesis of 1-benzyl-2,4,5-triphenyl-imidazole.

Figure 5: Synthesis of 1,2,4,5-tetrasubstituted imidazoles in the presence of nano-TiCl₄.SiO₂ at 110 °C under solvent-free condition.

($R^2 = 0.9998$). In this equation, A and C are the absorbance and concentration (mg L⁻¹) of titanium, respectively. The calculated loading amount of Ti in the catalyst is 7 mg g⁻¹.

The reaction of benzaldehyde (2 mmol), benzyl amine (2 mmol), benzil (1 mmol) and ammonium acetate (2 mmol) was examined to optimize the reaction conditions (Figure 4). Reaction under different conditions in the presence of TiCl₄.SiO₂ revealed that the use of 0.3 g of 50% TiCl₄.SiO₂, solvent-free, at 110 °C was the best condition (entry 16, Table 1). We repeated the above mentioned reaction, using 50% nano-TiCl₄.SiO₂, and found that 0.1 g of 50% nano-TiCl₄.SiO₂ was necessary (entry 19, Table 1). To examine the reusability of nano-TiCl₄.SiO₂, in a solvent-free condition, after each run, the product was dissolved to CHCl₃ and filtered. The catalyst residue was washed with acetone and reused. Treatment with acetone removes the tar from the catalyst surface more efficiently (Table 1, entries 22 and 23). The catalyst was reusable, although a gradual decline was observed in its activity.

Next, benzil, various aldehydes and amines were used as substrates for the synthesis of 1,2,4,5-tetra-substituted imidazoles (Figure 5 and Table 2). All products were known and characterized by FT-IR, ¹H-NMR, and the physical properties of those reported in the literature.

3. Conclusions

In conclusion, we have demonstrated a simple method for the synthesis of 1,2,4,5-tetra-substituted imidazoles using 50% nano-TiCl₄.SiO₂ as a reusable, eco-friendly, inexpensive and efficient reagent. Short reaction times, high yield, scale-up, simplicity of operation and easy work-up are some advantages of this method.

4. Experimental section

The chemicals were used without any additional purification. The products were characterized by FT-IR, ¹H-NMR, and a comparison of their physical properties with those reported in the literature. FT-IR spectra were run on a Bruker, Equinox 55 spectrometer. A Bruker (DRX-400 Avanes) NMR was used to record the ¹H NMR spectra. The absorption of titanium solutions was determined by an atomic absorption spectrometer; AVONA AA300. Also, the SEM of nano particles was determined with a VEGA/TESCAN scanning electron microscope.

4.1. Preparation of nano-TiCl₄.SiO₂ and TiCl₄.SiO₂

0.5 g (0.29 ml) of TiCl₄ was added drop wise to a mixture of 0.5 g of silica gel or nano-silica gel and 5 ml of chloroform.

Table 1: Acid catalyzed synthesis of 1-benzyl-2, 4, 5-triphenyl-imidazole.^a

Entry	Catalyst (g)	Solvent	Condition	Time (h)	Yield ^b (%)	Reference
1	TiCl ₄ .SiO ₂ (0.4)	Ethanol	r.t.	3	–	–
2	TiCl ₄ .SiO ₂ (0.4)	<i>n</i> -Hexane	r.t.	3	–	–
3	TiCl ₄ .SiO ₂ (0.4)	CHCl ₃	r.t.	3	10	–
4	TiCl ₄ .SiO ₂ (0.4)	CH ₂ Cl ₂	r.t.	3	15	–
5	TiCl ₄ .SiO ₂ (0.4)	ClCH ₂ CH ₂ Cl	r.t.	3	15	–
6	TiCl ₄ .SiO ₂ (0.4)	Ethanol	Reflux	2.5	30	–
7	TiCl ₄ .SiO ₂ (0.4)	<i>n</i> -Hexane	Reflux	2.5	30	–
8	TiCl ₄ .SiO ₂ (0.4)	CHCl ₃	Reflux	2.5	40	–
9	TiCl ₄ .SiO ₂ (0.4)	CH ₂ Cl ₂	Reflux	2.5	40	–
10	TiCl ₄ .SiO ₂ (0.4)	ClCH ₂ CH ₂ Cl	Reflux	2.5	45	–
11	TiCl ₄ .SiO ₂ (0.4)	Solvent-free	110 °C	2.5	75	–
12	30% TiCl ₄ .SiO ₂ (0.4)	Solvent-free	110 °C	2	70	–
13	50% TiCl ₄ .SiO ₂ (0.4)	Solvent-free	110 °C	1	85	–
14	60% TiCl ₄ .SiO ₂ (0.4)	Solvent-free	110 °C	1	85	–
15	50% TiCl ₄ .SiO ₂ (0.35)	Solvent-free	110 °C	1	84	–
16	50% TiCl ₄ .SiO ₂ (0.3)	Solvent-free	110 °C	1	84	–
17	50% TiCl ₄ .SiO ₂ (0.2)	Solvent-free	110 °C	1	75	–
18	50% Nano-TiCl ₄ .SiO ₂ (0.2)	Solvent-free	110 °C	0.5	95	–
19	50% Nano-TiCl ₄ .SiO ₂ (0.1)	Solvent-free	110 °C	0.5	94	–
20	50% Nano-TiCl ₄ .SiO ₂ (0.1)	Solvent-free	90 °C	0.5	88	–
21	50% Nano-TiCl ₄ .SiO ₂ (0.1)	Solvent-free	130 °C	0.5	95	–
22	50% Nano-TiCl ₄ .SiO ₂ (0.1), 2nd run	Solvent-free	110 °C	1	70	–
23	50% Nano-TiCl ₄ .SiO ₂ (0.1), 3rd run	Solvent-free	110 °C	1	58	–
24	I ₂ (5 mol%)	Ethanol	75 °C	15 min	95	[13]
25	K ₅ CoW ₁₂ O ₄₀ .3H ₂ O (0.1 mol%)	CH ₂ Cl ₂	140 °C	2	95	[14]
26	PEG-400 (1.5 g)	Solvent-free	110 °C	6	86	[21]
27	DABCO (0.7 mol%),	<i>t</i> -BuOH	60–65 °C	12–15	92	[20]
28	InCl ₃ .3H ₂ O (10 mol%)	Methanol	r.t.	8.3	82	[17]
29	37% BF ₃ /SiO ₂ (21 mol%)	Solvent-free	140 °C	2	92	[19]
30	SBPPSA (0.25 g, 0.31 mmol of H ⁺)	Solvent-free	140 °C	1	95	[22]

^a The used amount of benzaldehyde, benzyl amine, benzil and ammonium acetate is 2:2:1:2 mmol.^b Isolated yield.

The mixture was stirred for one hour at room temperature. The resulted suspension was filtered. The obtained solid was washed with chloroform and dried at room temperature.

4.2. General procedure for the synthesis of 1, 2, 4, 5-tetrasubstituted imidazoles

Benzil (1 mmol), amine (2 mmol), aldehyde (2 mmol), ammonium acetate (2 mmol) and 50% nano-TiCl₄.SiO₂ (0.1 g) were placed in a round bottom flask. The obtained mixture was stirred at 110 °C for 30 min. The progress of the reaction was followed by TLC. After completion of the reaction, the mixture was cooled to room temperature, dissolved in ethanol and filtered to separation of the catalyst. By addition of water to the concentrated filtrate, the solid product appeared. The product was re-crystallized in hot ethanol. All products are known and were identified by comparison of their physical and spectral data with those of authentic samples.

4.3. Selected spectroscopic data

1-Benzyl-2, 4, 5-triphenyl-imidazole: FT-IR(KBr) = 3020, 1598, 1490, 1435, 1382, 691, 754, ¹HNMR (400 MHz, CDCl₃): δ: 5.13 (s, 2H), 6.82 (sbr, 2H), 7.2 (m, 4H), 7.35 (m, 4H), 7.4 (m, 4H), 7.6 (d, *J* = 8 Hz, 2H), 7.67 (m, 4H) ppm.

1-Cyclohexyl-2, 4, 5-triphenyl-imidazole: FT-IR(KBr) = 3053, 2929, 2852, 1599, 1498, 1440, 1381, 1335, 745, 691, ¹HNMR (400 MHz, CDCl₃): δ: 0.8 (q br, *J* = 12 Hz, 1H), 1.1 (q br, *J* = 12 Hz, 2H), 1.6 (m, 6H), 1.9 (dbr, *J* = 12 Hz, 1H), 4 (t br, *J* = 12 Hz, 1H), 7.1 (m, 3H), 7.45 (m, 10H), 7.65 (m, 2H), ppm.

1-Benzyl-2-(4-chloro phenyl)-4, 5-diphenyl-imidazole: FT-IR (KBr) = 3100, 2900, 1601, 1497, 1448, 1028, 750, 696, ¹HNMR

Table 2: Nano-TiCl₄.SiO₂ catalyzed synthesis of 1, 2, 4, 5-tetrasubstituted imidazoles.^a

Entry	Ar	R	Yield (%) ^b	M.P. °C	Reference
1	C ₆ H ₅	C ₆ H ₅	82	217–218	[17]
2	C ₆ H ₅	C ₆ H ₅ CH ₂	84	162–165	[15]
3	C ₆ H ₅	Cyclohexyl	64	168–169	[12]
4	C ₆ H ₅	CH ₃ CH ₂	70	115–116	[12]
5	4-ClC ₆ H ₄	C ₆ H ₅	81	148–151	[14]
6	4-ClC ₆ H ₄	C ₆ H ₅ CH ₂	79	162–163	[17]
7	2-ClC ₆ H ₄	C ₆ H ₅ CH ₂	83	141–142	[17]
8	4-OHC ₆ H ₄	C ₆ H ₅ CH ₂	68	134–136	[17]
9	4-CH ₃ C ₆ H ₄	C ₆ H ₅	73	185–188	[16]
10	4-CH ₃ C ₆ H ₄	C ₆ H ₅ CH ₂	91	165–168	[12]
11	4-CH ₃ C ₆ H ₄	Cyclohexyl	75	163–164	[12]
12	3-OMeC ₆ H ₄	C ₆ H ₅ CH ₂	91	128–131	[17]
13	2-NO ₂ C ₆ H ₄	C ₆ H ₅ CH ₂	86	154–155	[17]
14	(CH ₃) ₂ CH	C ₆ H ₅ CH ₂	60	130–132	[17]

^a The ratio of benzil (mmol): aldehydes (mmol): amine (mmol): ammonium acetate (mmol): nano-TiCl₄.SiO₂ (g) is 1:2:2:2:0.1.^b Isolated yield.

(400 MHz, CDCl₃): δ: 5.1 (s, 2H), 6.8 (m, 2H), 7.25 (m, 7H), 7.4 (m, 6H), 7.6 (m, 4H), ppm.

2-(4-Chloro phenyl)-1, 4, 5-triphenyl-imidazole: FT-IR(KBr) = 3100, 2900, 1597, 1495, 1410, 1084, 836, 766, 696, ¹HNMR (400 MHz, CDCl₃): δ: 6.85 (d br, *J* = 6 Hz, 2H), 6.95 (d br, *J* = 6 Hz, 2H), 7.05 (m, 11H), 7.2 (d, *J* = 6 Hz, 2H), 7.4 (d, *J* = 6 Hz, 2H), ppm.

1-Benzyl, 2-(3-Methoxy phenyl)-4, 5-diphenyl-imidazole: FT-IR (KBr) = 3027, 2900, 1601, 1577, 1481, 1217, 695, ¹HNMR (400 MHz, CDCl₃): δ: 3.73 (s, 3H), 5.1 (s, 2H), 6.84 (m, 2H), 6.97 (m, 1H), 7.2 (m, 10H), 7.35 (m, 4H), 7.6 (d, *J* = 6 Hz, 2H), ppm.

1-Benzyl-2-(isopropyl)-4, 5-diphenyl-imidazole: FT-IR(KBr) = 3100, 2965, 1601, 1496, 1453, 1361, 1320, 750, 700, ¹HNMR (400 MHz, CDCl₃): δ: 1.2 (d, J = 6 Hz, 6H), 2.7 (m, 1H), 4.9 (s, 2H), 6.7 (d, J = 5.8 Hz, 2H), 7.1 (m, 11H), 7.3 (d, J = 5.8 Hz, 2H), ppm.

Acknowledgment

The Research Council of Yazd University is gratefully acknowledged for financial support of this study.

References

- [1] Laufer, S.A., Zimmermann, W. and Ruff, K.J. "Tetrasubstituted imidazole inhibitors of cytokine release: probing substituents in the N – 1 position", *J. Med. Chem.*, 47, pp. 6311–6325 (2004).
- [2] Misono, M. "Unique acid catalysis of heteropoly compounds (heteropoly-oxometalates) in the solid state", *Chem. Commun.*, pp. 1141–1152 (2001).
- [3] Black, J.W., Durant, G.J., Emmett, J.C. and Ganellin, C.R. "Sulphurmethyle isosterism in the development of metiamide, a new histamine H₂-receptor antagonist", *Nature*, 248, pp. 65–67 (1974).
- [4] Ucuu, U., Karaburun, N.G. and Iskdag, I. "Synthesis and analgesic activity of some 1-benzyl-2-substituted-4, 5-diphenyl-1H-imidazole derivatives", *Il Farmaco*, 56, pp. 285–290 (2001).
- [5] Murry, J.A. "Synthetic methodology utilized to prepare substituted imidazole p38MAPkinase inhibitors. Curr Opin", *Drug Discov. Dev.*, 6, pp. 945–965 (2003).
- [6] de Laszlo, S.E., Hacker, C., Li, B., Kim, D., MacCoss, M., Mantlo, N., Pivnichny, J.V., Colwell, L., Koch, G.E., Cascieri, M.A. and Hagmann, W.K. "Potent, orally absorbed glucagon receptor antagonists", *Bioorg. Med. Chem. Lett.*, 9, pp. 641–646 (1999).
- [7] Eysers, P.A., Craxton, M., Morrice, N., Cohen, P. and Goedert, M. "Conversion of SB 203580-insensitive MAP kinase family members to drug-sensitive forms by a single amino-acid substitution", *Chem. Biol.*, 5, pp. 321–328 (1998).
- [8] Newman, M.J., Rodarte, J.C., Benbatoul, K.D., Romano, S.J., Zhang, C., Krane, S., Moran, E.J., Uyeda, R.T., Dixon, R., Guns, E.S. and Mayer, L.D. "Discovery and characterization of OC144-093, a novel inhibitor of P-glycoprotein-mediated multidrug resistance", *Cancer Res.*, 60, pp. 2964–2972 (2000).
- [9] Antolini, M., Bozzoli, A., Ghiron, C., Kennedy, G., Rossi, T. and Ursini, A. "Analogues of 4, 5-bis(3, 5-dichlorophenyl)-2-trifluoromethyl-1H-imidazole as potential antibacterial agents", *Bioorg. Med. Chem. Lett.*, 9, pp. 1023–1028 (1999).
- [10] Wang, L., Woods, K.W., Li, Q., Barr, K.J., McCroskey, R.W., Hannick, S.M., Gherke, L., Credo, R.B., Hui, Y.-H., Marsh, K., Warner, R., Lee, J.Y., Zielinsky-Mozng, N., Frost, D., Rosenberg, S.H. and Sham, H.L. "Potent, orally active heterocycle-based combretastatin α-4 analogues: synthesis, structure-activity relationship, pharmacokinetics, and in vivo antitumor activity evaluation", *J. Med. Chem.*, 45, pp. 1697–1711 (2001).
- [11] Balalaei, S. and Arabanian, A. "One-pot synthesis of tetrasubstituted imidazoles catalyzed by zeolite HY and silica gel under microwave irradiation", *Green Chem.*, 2, pp. 274–276 (2000).
- [12] Karimi, A.R., Alimohammadi, Z., Azizian, J., Mohammadi, A.A. and Mohmmadizadeh, M.R. "Solvent-free synthesis of tetrasubstituted imidazoles on silica gel/NaHSO₄ support", *Catal. Commun.*, 7, pp. 728–732 (2006).
- [13] Kidwai, M., Mothsra, P., Bansal, V., Somvanshi, R.K., Ethayathulla, A.S., Dey, S. and Singh, T.P. "One-pot synthesis of highly substituted imidazoles using molecular iodine: a versatile catalyst", *J. Mol. Catal. A: Chem.*, 265, pp. 177–182 (2007).
- [14] Nagarapu, L., Apuri, S. and Kantevari, S. "Potassium dodecatungstocobaltate trihydrate (K₅CoW₁₂O₄₀·3H₂O): a mild and efficient reusable catalyst for the one-pot synthesis of 1, 2, 4, 5-tetrasubstituted imidazoles under conventional heating and microwave irradiation", *J. Mol. Catal. A: Chem.*, 266, pp. 104–108 (2007).
- [15] Heravi, M.M., Derikvand, F. and Bamoharram, F.F. "Highly efficient, four-component one-pot synthesis of tetrasubstituted imidazoles using Keggin-type heteropolyacids as green and reusable catalysts", *J. Mol. Catal. A: Chem.*, 263, pp. 112–114 (2007).
- [16] Kantevari, S., Vuppapapati, S.V.N., Biradar, D.O. and Nagarapu, L. "Synthesis of 1, 2, 4, 5-tetrasubstituted imidazoles using silica-bonded propylpiperazine N-sulfamic acid as a recyclable solid acid catalyst", *J. Mol. Catal. A: Chem.*, 266, pp. 109–113 (2007).
- [17] Sharma, S.D., Hazarika, P. and Konwar, D. "An efficient and one-pot synthesis of 2, 4, 5-trisubstituted and 1, 2, 4, 5-tetrasubstituted imidazoles catalyzed by InCl₃·3H₂O", *Tetrahedron Lett.*, 49, pp. 2216–2220 (2008).
- [18] Sharma, G.V., Jyothi, Y. and Lakshmi, P.S. "Efficient room-temperature synthesis of tri- and tetrasubstituted imidazoles catalyzed by ZrCl₄", *Synth. Commun.*, 36, pp. 2991–3000 (2006).
- [19] Sadeghi, B., Mirjalili, B.F. and Hashemi, M.M. "BF₃·SiO₂ an efficient reagent system for the one-pot synthesis of 1, 2, 4, 5-tetrasubstituted imidazoles", *Tetrahedron Lett.*, 49, pp. 2575–2577 (2008).
- [20] Murthy, S.N., Madhav, B. and Nageswar, Y.V.D. "DABCO as a mild and efficient catalytic system for the synthesis of highly substituted imidazoles via multi-component condensation strategy", *Tetrahedron Lett.*, 51, pp. 5252–5257 (2010).
- [21] Wang, X.C., Gong, H.P., Quan, Z.J., Li, L. and Ye, H.L. "PEG-400 as an efficient reaction medium for the synthesis of 2, 4, 5-triaryl-1H-imidazoles and 1, 2, 4, 5-tetraaryl-1H-imidazoles", *Chinese Chem. Lett.*, 20, pp. 44–47 (2009).
- [22] Niknam, K., Deris, A., Naeimi, F. and Majleci, F. "Synthesis of 1,2,4,5-tetrasubstituted imidazoles using silica-bonded propylpiperazine N-sulfamic acid as a recyclable solid acid catalyst", *Tetrahedron Lett.*, 52, pp. 4642–4645 (2011).
- [23] Souza, J.L.D., Fabri, F., Buffon, R. and Schuchardt, R. "Reduced anatase on silica as a support for a Ziegler catalyst", *Appl. Catal. A. Gen.*, 323, p. 234 (2007).
- [24] Mirjalili, B.F., Bamoniri, A. and Akbari, A. "BF₃·SiO₂: an efficient alternative for the synthesis of 14-aryl or alkyl-14H-dibenzo[a, j]xanthenes", *Tetrahedron Lett.*, 49, pp. 6454–6456 (2008).
- [25] Sadeghi, B., Mirjalili, B.F. and Hashemi, M.M. "BF₃·SiO₂: an efficient heterogeneous alternative for regio-chemo and stereoselective claisen-schmidt condensation", *J. Iran. Chem. Soc.*, 8, pp. 694–698 (2011).
- [26] Mirjalili, B.F., Bamoniri, A. and Akbari, A. "One-pot synthesis of 3, 4-dihydropyrimidin-2(1H)-ones(thiones) promoted by nano-BF₃·SiO₂", *J. Iran. Chem. Soc.*, 8, pp. S135–S140 (2011).
- [27] Zettler, V.A. and Brown, C.A. "Infrared spectra of some Ti–O–Si, Ti–O–Ti and Si–O–Si compounds", *J. Am. Chem. Soc.*, 79, pp. 1174–1178 (1957).
- [28] Schrijnemakers, K., Voort, P.V.D. and Vansant, E.F. "Characterization of a silica surface TiCl₄-modified by means of quantitative surface analysis", *Phys. Chem. Chem. Phys.*, 1, pp. 2569–2572 (2008).

Bi Bi Fatemeh Mirjalili was born in Yazd, Iran, in 1961. She obtained a B.S. degree in Chemistry from Alzahra University, Tehran, Iran, in 1987, an M.S. degree in Organic Chemistry from Tarbiat Moalem University, Iran, in 1991, and a Ph.D. degree in Organic Chemistry from Sharif University of Technology, Tehran, Iran, in 2001.

She has been Full Professor at Yazd University, Iran, since 2011.

Abdol Hamid Bamoniri was born in Abadan, Iran, in 1958. He obtained a B.S. degree in Chemistry from Shahid Beheshti University, Tehran, Iran, in 1984, an M.S. degree in Organic Chemistry from Tarbiat Moalem University, Tehran, Iran, in 1989, and a Ph.D. degree in Organic Chemistry from Bu-Ali Sina University, Hamedan, Iran, in 2003.

He has been Associate Professor at Kashan University, Iran, since 2007.

Leila Zamani was born in Abadeh, Iran, in 1983. She obtained her B.S. and M.S. degrees in Chemistry and Organic Chemistry from Yazd University, Iran, in 2005 and 2008, respectively, where she is now a Ph.D. degree student in the same subject.